

## DCIS – QUESTIONS

Clinical Case Conference  
UCSD Radiation Oncology  
SA-CME

1. To be included in the low risk group in the ECOG DCIS trial a patient could have all of the following except?
  - A) Grade 2 Disease
  - B) 2.9 cm lesion
  - C) ER positive disease
  - D) 3mm margin
  - E) Non-palpable lesion
2. Patients enrolled in the UK Tamoxifen trial who did not get radiotherapy and did not get tamoxifen had a 10% ipsilateral recurrence rate of DCIS. What was the recurrence rate for patients that were randomized to tamoxifen only?
  - A) 10%
  - B) 8%
  - C) 6%
  - D) 4%
  - E) 2%
3. What percentage of women discontinued treatment (tamoxifen or placebo) in the NSABP B-24 trial?
  - A) 61%
  - B) 51%
  - C) 41%
  - D) 31%
  - E) 21%
4. Which of the following risk factors is not included in the USC/Van Nuys prognostic index (USC/VNPI)?
  - A) Tumor size
  - B) Margin width
  - C) Age
  - D) Histologic classification
  - E) Hormone receptor status
5. NSABP B-43 is a trial examining the role of trastuzumab in the treatment of women with Her2neu+ DCIS. Approximately what percentage women enrolled on the trial were Her2neu+?
  - A) 25%
  - B) 35%
  - C) 45%
  - D) 55%
  - E) 65%